## **BIOPRIME - CHILLI CROP SCHEDULE**











- Helps in additional picking of the fruits.



	TRICHONEXUS	O. J. WERL	DH4	PRII	P P P P P P P P P P P P P P P P P P P
Basal Dose	Vegetative Stage ( 2 - 5 Days )	Vegetative Stage ( 10 - 20 Days )	Vegetative Stage ( 20 - 25 Days )	Flowering Stage ( 40 - 45 Days )	Fruits Development & Maturity ( 55 – 60 Days )
Prime CG - Pro (10 Kg / Acre)	Prime 1515 (1000 ml) Prime Trichonexus (1 kg)	Prime Verdant (500 ml) Prime Fortisea (500 ml)	Phosphonexus (1000 ml)	Prime 7525 (500 ml) Blossom k nexus (1000 ml)	Prime Admiron (500 ml) Prime Fortisea (1000 ml)
<ul> <li>Promotes growth of white roots.</li> <li>Increases bioavailability of fertilizers.</li> <li>Helps to keep soil Porus.</li> <li>Mitigates transplantation shock &amp; promotes establishment.</li> </ul>		<ul> <li>Confers resistance against abiotic stress, mitigates biotic stress pressure.</li> <li>Accelerates the photosynthetic activity.</li> <li>Promotes white root growth, enhances uptake of phosphorus.</li> </ul>		<ul> <li>Reduces flower droping.</li> <li>Increases flower to fruit conversion.</li> <li>Enhances fruit weight &amp; length.</li> <li>Uplifts fruit quality &amp; shine.</li> <li>Increased shelf life of fruits.</li> <li>Enhances uptake of potassium nutrients.</li> </ul>	

